


BI



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) EP 0 756 869 A2

(12) EUROPEAN PATENT APPLICATION

(43) Date of publication:
05.02.1997 Bulletin 1997/06

(51) Int Cl⁶ A61K 31/135, A61K 31/38

(21) Application number: 96305346.7

(22) Date of filing: 22.07.1996

(84) Designated Contracting States
AT BE CH DE DK ES FI FR GB GR IE IT LI LU NL
PT SE
Designated Extension States
AL LT LV SI

(30) Priority: 24.07.1995 US 1373
08.11.1995 US 6349
19.12.1995 GB 9525909

(71) Applicant: ELI LILLY AND COMPANY
Indianapolis, Indiana 46285 (US)

(72) Inventors:
• Heiligenstein, John Harrison
Indianapolis, Indiana 46228 (US)

• Laguzza, Bennett Coleman
Indianapolis, Indiana 46250 (US)
• Paul, Steven Marc
Carmel, Indiana 46032 (US)
• Tollefson, Gary Dennis
Indianapolis, Indiana 46236 (US)

(74) Representative: Hudson, Christopher Mark
Lilly Industries Limited
European Patent Operations
Erl Wood Manor
Windlesham Surrey GU20 6PH (GB)

(54) Use of duloxetine or of N-alkyl-3-phenyl-(2-alkylthiophenoxy)-propylamines for treating attention-deficit/hyperactivity disorder

(57) A group of propanamines, norepinephrine up-

take inhibitors, are used to treat attention-deficit/hyperactivity disorder.

EP 0 756 869 A2

Description

The invention belongs to the fields of pharmaceutical chemistry and psychiatric medicine, and provides a method of treatment of the psychiatric disorder known as attention-deficit/hyperactivity disorder


For some decades it has been recognized that a significant number of children are persistently hyperactive and have an attention span so short as to be disabling in school and in many personal relationships. Such children in earlier times would no doubt have been dismissed as incorrigible and punished or even confined in an institution. Some long time ago, however, it was realized that these children cannot control their hyperactivity and inattention, and the medical professions began to try to help them. Methylphenidate (RitalinTM) has been used for some time to treat such children and it often significantly improves their ability to function and coexist with other people at school and at home. However, the drug has the disadvantages of requiring several doses per day, and producing a rebound effect as the effect of each dose fades away. Further, the drug causes sleeplessness and lack of appetite in some patients. Methylphenidate has both noradrenergic and dopaminergic activities.

Imipramine, desipramine, nortriptyline, amitriptyline and clomipramine are also used in some cases of attention-deficit/hyperactivity disorder (ADHD). Those tricyclic drugs, however, have a number of physiological mechanisms and, as a class, tend to produce a number of side effects and require careful supervision and dose titration.

In the last decade, psychiatrists have realized that ADHD is not only a disorder of childhood, but often continues in the adult. It is obvious that hyperactivity and short attention span cause grave disruption in an adult's life, but it is only recently that such patients have been able to obtain any treatment.

The need for a safe and convenient treatment for ADHD, applicable to both children and adults and without the disadvantages possessed by methylphenidate continues to be a concern of the psychiatric profession.

The present invention provides the use for manufacturing a medicament for the treatment of attention-deficit/hyperactivity disorder of duloxetine, or of a compound of the formula



wherein X = C₁-C₄ alkylthio, and Y = C₁-C₂ alkyl or a stereoisomer thereof, or a pharmaceutically acceptable salt thereof.

Duloxetine, N-methyl-3-(1-naphthalenyloxy)-3-(2-thienyl)propanamine, is usually administered as the hydrochloride salt and as the (+) enantiomer. It was first taught by U.S. Patent 4,956,388, which shows its high potency. The word "duloxetine" will be used here to refer to any acid addition salt or the free base of the molecule.

The compounds of formula I were described in U.S. Patent 5,281,624, of Gehlert, Robertson and Wong, and in Gehlert, et al., *Life Sciences*, 56(22), 1915-1920, 1995. The compounds are there taught to be inhibitors of norepinephrine reuptake in the brain. It is also explained that the compounds exist as stereoisomers, and that they accordingly include not only the racemates, but also the isolated individual isomers. For example, the compounds of formula I include the following exemplary species

N-ethyl-3-phenyl-3-(2-methylthiophenoxy)propylamine benzoate
 (R)-N-methyl-3-phenyl-3-(2-propylthiophenoxy)-propylamine hydrochloride
 (S)-N-ethyl-3-phenyl-3-(2-butylthiophenoxy)-propylamine
 N-methyl-3-phenyl-3-(2-ethylthiophenoxy)propylamine malonate
 (S)-N-methyl-3-phenyl-3-(2-1-butylthiophenoxy)-propylamine naphthalene-2-sulfonate
 (R)-N-methyl-3-(2-methylthiophenoxy)-3-phenylpropylamine
 (S)-N-methyl-3-phenyl-3-(2-1-butylthiophenoxy)-propylamine naphthalene-2-sulfonate

The compounds used in this invention are safe drugs, and their use in ADHD, in both adults and children, is a superior treatment for that disorder because of their improved safety. The compounds are particularly selective, having few if any physiological effects besides that on norepinephrine processing, and therefore are free of side effects and unwanted activities. Further, they are effective at relatively low doses as discussed below, and may safely and effec-

tively be administered once per day. Thus, difficulties created by the multiple dosing of patients, who are children and disorganized adults, are completely avoided.

The present invention is described as a method of treating ADHD. It is not intended, of course, that every use of the invention will result in a complete cure of the disorder. Treatment of psychiatric disorders is not that precise. It is intended, however, that the use of the invention will result in improvement of the treated ADHD patient which will approach complete cure in some cases. In other cases, a lesser degree of improvement, still benefiting the patient, will be obtained. Such treatment is carried out by administering an effective amount of the chosen compound. It will be understood that amount is an amount which produces treatment of the patient at hand.

The effective dose of a compound of this invention for ADHD is in the range from about 1 mg/day to about 100 mg/day. The preferred adult dose is in the range from about 5 to about 80 mg/day, and a more highly preferred adult dose is from about 10 to about 60 mg/day. The children's dose of course is smaller, in the range from about 1 to about 70 mg/day, more preferably from about 5 to about 60 mg/day and still more preferably from about 5 to about 50 mg/day. The optimum dose for each patient, as always, must be set by the physician in charge of the case, taking into account the patient's size, other medications which the patient requires, severity of the disorder and all of the other circumstances of the patient.

Since the compounds are readily orally absorbed and require only once/day administration, there is little or no reason to administer them in any other way than orally. They may be produced in the form of a clean, stable crystal, and thus are easily formulated in the usual oral pharmaceutical forms, such as tablets, capsules, suspensions and the like. The usual methods of pharmaceutical scientists are applicable. They may usefully be administered, if there is any reason to do so in a particular circumstance, in other pharmaceutical forms; such as injectable solutions, depot injections, suppositories and the like, which are well known to and understood by pharmaceutical scientists. It will substantially always be preferred, however, to administer a compound as a tablet or capsule and such pharmaceutical forms are recommended.

A preferred duloxetine enteric formulation is a pellet formulation comprising a) a core consisting of duloxetine and a pharmaceutically acceptable excipient; b) an optional separating layer; c) an enteric layer comprising hydroxypropylmethylcellulose acetate succinate (HPMCAS) and a pharmaceutically acceptable excipient; d) an optional finishing layer. The following example demonstrates the preparation of a preferred such formulation.

Example

10 mg Duloxetine base/capsule

Bill of Materials

Beads

Sucrose - starch nonpareils,

20-25 mesh 60.28 mg

Duloxetine layer

Duloxetine 11.21

Hydroxypropylmethylcellulose 3.74

Separating layer

Hydroxypropylmethylcellulose 2.51

Sucrose 5.00

Talc, 500 mesh 10.03

Enteric layer

HPMCAS, LF grade, Shin-Etsu Chemical 25.05

Co., Tokyo, Japan

Triethyl citrate 5.00

Talc, 500 mesh 7.52

Finishing layer

Hydroxypropylmethylcellulose 8.44

Titanium dioxide 2.81

Talc Trace

141.60 mg

The duloxetine layer was built up by suspending duloxetine in a 4% w/w solution of the hydroxypropylmethylcellulose in water, and milling the suspension with a CoBall Mill (Fryma Mashinen AG, Rheinfelden, Switzerland) model MS-12. A fluid bed dryer with a Wurster column was used to make this product, at a batch size of 10 kg. The separating layer was added from a 4% w/w solution of the hydroxypropylmethylcellulose in water, in which the sucrose was also dissolved.

In order to prepare the enteric coating suspension, purified water was cooled to 10°C and the polysorbate, triethyl citrate and silicone emulsion were added and dispersed or dissolved. Then the HPMCAS and talc were added and agitated until homogeneity was obtained, and the HPMCAS was fully neutralized by addition of ammonium hydroxide until solution of the polymer was complete. To this suspension, a carboxymethylcellulose aqueous solution, 0.5% w/w, was added and blended thoroughly. The enteric suspension was maintained at 20°C during the coating process. The enteric suspension was then added to the partially completed pellets in the Wurster column at a spray rate of about 15 ml/min, holding the temperature of the inlet air at about 50°C. The product was dried in the Wurster at 50°C when the enteric suspension had been fully added, and then dried on trays for 3 hours in a dry house at 60°C. A finishing layer was then applied which consisted of a 4.5% w/w hydroxypropylmethylcellulose solution containing titanium dioxide and propylene glycol as plasticizer. The pellets were completely dried in the fluid bed dryer and then were then filled in size 3 gelatin capsules.

The ADHD patient is rather readily recognized, and most people have been in contact with children, if not adults, who exhibit some or all of the symptoms of the disorder. The best description of the disorder is the diagnostic criteria for ADHD published by the American Psychiatric Association in the Diagnostic and Statistical Manual of Mental Disorders, Fourth Version (1994), as follows

Diagnostic criteria for Attention-Deficit/Hyperactivity Disorder

A Either (1) or (2)

(1) six (or more) of the following symptoms of **inattention** have persisted for at least 6 months to a degree that is maladaptive and inconsistent with developmental level

Inattention

- (a) often fails to give close attention to details or makes careless mistakes in schoolwork, work, or other activities
- (b) often has difficulty sustaining attention in tasks or play activities
- (c) often does not seem to listen when spoken to directly
- (d) often does not follow through on instructions and fails to finish schoolwork, chores, or duties in the workplace (not due to oppositional behavior or failure to understand instructions)
- (e) often has difficulty organizing tasks and activities
- (f) often avoids, dislikes, or is reluctant to engage in tasks that require sustained mental effort (such as schoolwork or homework)
- (g) often loses things necessary for tasks or activities (e.g., toys, school assignments, pencils, books, or tools)
- (h) is often easily distracted by extraneous stimuli
- (i) is often forgetful in daily activities

(2) six (or more) of the following symptoms of **hyperactivity-impulsivity** have persisted for at least 6 months to a degree that is maladaptive and inconsistent with developmental level.

Hyperactivity

- (a) often fidgets with hands or feet or squirms in seat
- (b) often leaves seat in classroom or in other situations in which remaining seated is expected
- (c) often runs about or climbs excessively in situations in which it is inappropriate (in adolescents or adults, may be limited to subjective feelings of restlessness)
- (d) often has difficulty playing or engaging in leisure activities quietly
- (e) is often "on the go" or often acts as if "driven by a motor"
- (f) often talks excessively *Impulsivity*
- (g) often blurts out answers before questions have been completed
- (h) often has difficulty awaiting turn
- (i) often interrupts or intrudes on others (e.g., butts into conversations or games)

B Some hyperactive-impulsive or inattentive symptoms that caused impairment were present before age 7 years

C Some impairment from the symptoms is present in two or more settings (e.g., at school [or work] and at home)

D There must be clear evidence of clinically significant impairment in social, academic, or occupational functioning

E The symptoms do not occur exclusively during the course of a Pervasive Developmental Disorder, Schizophrenia, or other Psychotic Disorder and are not better accounted for by another mental disorder (e.g., Mood Disorder, Anxiety Disorder, Dissociative Disorder, or a Personality Disorder)

It will be seen that ADHD is a disorder made up of two components, the attention deficit component and the hyperactivity component, which are to a degree independent. Treatment with the compounds is effective in patients who are primarily suffering from either component or from the combined disorder.

While ADHD is still primarily regarded as a disorder of children, it is now understood that many ADHD patients, as many as 50% continue to suffer from the disorder as they grow through adolescence into adulthood. Biederman and associates have extensively studied the adult ADHD patient, and have found numerous cases. See, for example, Biederman, et al. *Am J Psychiatry* 150, 1792-98 (1993). They found that cases of adult ADHD were frequently found among the parents and adult siblings of childhood ADHD patients. Thus, it appears that the disease is not only carried forward into adulthood, but is inheritable.


The Biederman, et al. article cited immediately above, as well as another article by the same authors, *Am J Psychiatry* 148, 564-77 (1991), reports studies of ADHD patients who also have one or more other psychiatric disorders. The authors indicate that such psychiatric co-morbidity is quite common among ADHD patients and, naturally, cloud the diagnosis and treatment of such patients. The compounds are effective in the treatment of ADHD, even though the situation of the treated patient may be complicated by co-morbidity with one or more additional disorders.

The mere listing of the above diagnostic criteria indicates the seriousness of ADHD and the damage which it does to the patient. A person having a moderately severe case of ADHD is substantially entirely unable to concentrate and hence unable to do meaningful work or study, is a continuing distraction and nuisance to those around her or him, because of the uselessly impulsive activity which the disorder causes, and consumes his or her family in cleaning up and repairing the damage and disruption which he or she causes. Such a patient of school age may substantially damage the teacher's ability to accomplish the class' goals, because the ADHD child will continually disrupt the class, distract the other children and consume the teacher's effort. Thus, it is readily apparent that an improved treatment of ADHD is needed, and that the present invention is accordingly important to many people.

The method of the present invention is effective in the treatment of patients who are children, adolescents or adults, and there is no significant difference in the symptoms or the details of the manner of treatment among patients of different ages. In general terms, however, for purposes of the present invention, a child is considered to be a patient below the age of puberty, an adolescent is considered to be a patient from the age of puberty up to about 18 years of age, and an adult is considered to be a patient of 18 years or older.

Claims


1. The use for the manufacture of a medicament for treating attention-deficit/hyperactivity disorder of duloxetine, or of a compound of the formula



wherein X = C₁-C₄ alkylthio, and Y = C₁-C₂ alkyl or a stereoisomer thereof, or a pharmaceutically acceptable salt thereof.

2. A use of Claim 1 wherein the predominantly inattentive type of attention-deficit/hyperactivity disorder is to be treated.
3. A use of Claim 1 wherein the predominantly hyperactive-impulsive type of attention-deficit/hyperactivity disorder is to be treated.
4. A use of Claim 1 wherein the combined type of attention-deficit/hyperactivity disorder is to be treated.
5. A use of Claim 1 wherein the patient to be treated is a child.
6. A use of Claim 1 wherein the patient to be treated is an adolescent.
7. A use of Claim 1 wherein the patient to be treated is an adult.
8. A use of any one of Claims 1-7 wherein the compound is duloxetine.
9. A use of any one of Claims 1-7 wherein the compound is N-methyl-3-(2-methylthiophenoxy)-3-phenylpropylamine or a stereoisomer thereof, or a pharmaceutically acceptable salt thereof.

Best Available Copy



(19)

Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 0 756 869 A3

(12)

EUROPEAN PATENT APPLICATION

(88) Date of publication A3
16.06.1999 Bulletin 1999/24

(51) Int Cl⁶ **A61K 31/135, A61K 31/38**

(43) Date of publication A2
05.02.1997 Bulletin 1997/06

(21) Application number: **96305346.7**

(22) Date of filing: **22.07.1996**

(84) Designated Contracting States:
**AT BE CH DE DK ES FI FR GB GR IE IT LI LU NL
PT SE**
Designated Extension States:
AL LT LV SI

(30) Priority **24.07.1995 US 1373 P**
08.11.1995 US 6349 P
19.12.1995 GB 9525909

(71) Applicant **ELI LILLY AND COMPANY**
Indianapolis, Indiana 46285 (US)

(72) Inventors
• Heiligenstein, John Harrison
Indianapolis, Indiana 46228 (US)

• Laguzza, Bennett Coleman
Indianapolis, Indiana 46250 (US)
• Paul, Steven Marc
Carmel, Indiana 46032 (US)
• Tollefson, Gary Dennis
Indianapolis, Indiana 46236 (US)

(74) Representative **Hudson, Christopher Mark**
Lilly Industries Limited
European Patent Operations
Erl Wood Manor
Windlesham Surrey GU20 6PH (GB)

(54) **Use of duloxetine or of N-alkyl-3-phenyl-(2-alkylthiophenoxy)-propylamines for treating attention-deficit/hyperactivity disorder**

(57) A group of propanamines, norepinephrine up-

take inhibitors, are used to treat attention-deficit/hyperactivity disorder.

EP 0 756 869 A3

Best Available Copy



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 96 30 5346

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
X	ZA 930 694 A (ELI LILLY) * page 49, paragraph 2 * * page 46, last paragraph * ---	1-8	A61K31/135 A61K31/38
P,X	EP 0 687 472 A (ELI LILLY) 20 December 1995 * page 14, line 25; claims 1,3,6,7,10,11 * * page 14, line 8-12 * ---	1-8	
P,X	WO 96 12485 A (ELI LILLY) 2 May 1996 * page 9, column 3-15; claims 1-5 * * page 5, line 20 * ---	1-8	
Y	US 5 114 976 A (NORDEN MICHAEL J) 19 May 1992 * tables 1-CASE4 * ---	1-8	
D,Y	US 4 956 388 A (ROBERTSON DAVID W ET AL) 11 September 1990 * column 1; example 2 * ---	1-8	
Y	EP 0 537 915 A (ELI LILLY) 21 April 1993 * page 2, line 11-45; claims * D & US 5 281 624 A ---	1-7,9	TECHNICAL FIELDS SEARCHED (Int.Cl.6) A61K
Y	EP 0 591 581 A (ELI LILLY) 13 April 1994 * page 2, line 15-47 * ---	1-7,9	
D,Y	D.R. GEHLERT: "(R)-Thionisoxetine, a potent and selective inhibitor of central and peripheral norepinephrine uptake" LIFE SCIENCES, vol. 56, no. 22, April 1995, pages 1915-1920, XP002092329 * last paragraph * ---	1-7,9	
-/--			
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 4 February 1999	Examiner ORVIZ DIAZ P.
CATEGORY OF CITED DOCUMENTS X particularly relevant if taken alone Y particularly relevant if combined with another document of the same category A technological background C non-written disclosure P intermediate document		T theory or principle underlying the invention E earlier patent document, but published on, or after the filing date D document cited in the application L document cited for other reasons & member of the same patent family, corresponding document	



European Patent
Office

Application Number

EP 96 30 5346

CLAIMS INCURRING FEES

The present European patent application comprised at the time of filing more than ten claims.

- ☐ Only part of the claims have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims and for those claims for which claims fees have been paid, namely claim(s):
- ☐ No claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims.

LACK OF UNITY OF INVENTION

The Search Division considers that the present European patent application does not comply with the requirements of unity of invention and relates to several inventions or groups of inventions, namely:

see sheet B

- ☒ All further search fees have been paid within the fixed time limit. The present European search report has been drawn up for all claims.
- ☐ Only part of the further search fees have been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the inventions in respect of which search fees have been paid, namely claims:
- ☐ None of the further search fees have been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the invention first mentioned in the claims, namely claims:

BEST AVAILABLE COPY

European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 96 30 5346

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (InCL6)
T	US 5 696 168 A (J.H. HEILIGENSTEIN) 9 December 1997 * the whole document *	1-7,9	
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 4 February 1999	Examiner ORVIZ DIAZ P.
CATEGORY OF CITED DOCUMENTS		T theory or principle underlying the invention E earlier patent document, but published on, or after the filing date D document cited in the application L document cited for other reasons A technological background O non-written disclosure P intermediate document & member of the same patent family, corresponding document	
X particularly relevant if taken alone Y particularly relevant if combined with another document of the same category			

EPO FORM 1503 01/92 (P01C01)

BEST AVAILABLE COPY

European Patent
OfficeLACK OF UNITY OF INVENTION
SHEET BApplication Number
EP 96 30 5346

The Search Division considers that the present European patent application does not comply with the requirements of unity of invention and relates to several inventions or groups of inventions, namely:

1. Claims: 1-7 (partially) and 8 (complete)

Use of duloxetine for the manufacture of a medicament for treating attention-deficit/hyperactivity disorder (ADHD)

2. Claims: 1-7 (partially) and 9 (complete)

Use of N-alkyl-3-phenyl-3-(2-alkylthiophenoxy)propylamines of the formula in claim 1 for the manufacture of a medicament for treating attention-deficit/hyperactivity disorder (ADHD)